

7114-B1

BOVA-CREEP MEDICATED

FOR BEEF CATTLE ON PASTURE

For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle, and beef replacement heifers.)

ACTIVE DRUG INGREDIENT

BAMBERMYCINS 4 gm/ton

NO WITHDRAWAL PERIOD REQUIRED

GUARANTEED ANALYSIS

CRUDE PROTEIN	MIN	14.00 %	SALT SALT POTASSIUM SELENIUM	MIN	1.00 %
CRUDE FAT	MIN	3.50 %	SAIT	NAAV	2.00 %
CRUDE FIBER	MAX	13.50 %	DOTASSILIM	NAIN	0.6 %
CALCIUM	MIN	1.75 %	CELENILIM	DAIN	0.8 % 0.30 PPM
CALCIUM	MAX	2.25 %	VITAMIN A	IVIIIN	0.30 PPIVI
PHOSPHORUS	MIN	0.50 %	VITAWIIN A	IVIIIN	4000 IU/LB

INGREDIENTS

PROCESSED GRAIN BY-PRODUCTS, PLANT PROTEIN PRODUCTS, GRAIN PRODUCTS, MOLASSES PRODUCTS GROUND LIMESTONE, ROUGHAGE PRODUCTS (15%), SALT, VITAMIN A SUPPLEMENT, VITAMIN D3 SUPPLEMENT, VITAMIN E SUPPLEMENT, ZINC PROTEINATE, COPPER PROTEINATE, MANGANESE PROTEINATE, COBALT PROTEINATE, ZINC SULFATE, CALCIUM CARBONATE, FERROUS CARBONATE, MANGANESE SULFATE, COPPER SULFATE, SODIUM SELENITE, ETHYLENEDIAMINE DIHYDRIODIDE, COBALT SULFATE, DRIED ASPERGILLUS ORYZAE FERMENTATION PRODUCT, HYDROLYZED YEAST, BREWERS DRIED YEAST, DRIED BACILLUS LICHENIFORMIS FERMENTATION PRODUCT, DRIED BACILLUS SUBTILIS FERMENTATION PRODUCT.

FEEDING DIRECTIONS

FEED CONTIUNALLY AT A RATE OF 10-40 MG/HEAD/DAY IN A LEAST 1 POUND AND NOT MORE THAN 10 POUNDS OF FEED. DAILY BAMBERMYCINS INTAKE IN EXCESS OF 20 MG/HEAD/DAY HAVE NOT BEEN SHOWN TO BE MORE EFFECTIVE THAN 20 MG/HEAD/DAY.

CAUTION: CONTAINS COPPER: COPPER HAS BEEN SHOWN TO BE TOXIC TO SHEEP.

3/8

ALLIED FEEDS INC. 208 HUTCHESON, CUERO, TEXAS 77954 1-800-242-3500 NET WT 50# (22.72 KG) 01

