

7114-B1

BOVA-CREEP MEDICATED

FOR BEEF CATTLE ON PASTURE

For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle, and beef replacement heifers.)

ACTIVE DRUG INGREDIENT

BAMBERMYCINS 4 gm/ton

NO WITHDRAWAL PERIOD REQUIRED

GUARANTEED ANALYSIS

CRUDE PROTEIN	MIN	14.00 %	SALT SALT POTASSIUM SELENIUM	MIN	1 00 %
CRUDE FAT	MIN	3.50 %	SAIT	NAAV	2.00 %
CRUDE FIBER	MAX	13.50 %	DOTASSILIM	NAIN	2.00 %
CALCIUM	MIN	1.75 %	SELENILIM	NAIN	0.0 %
CALCIUM	MAX	2.25 %	VITAMINI A	IVIIIN	0.30 PPIVI
PHOSPHORUS	MIN	0.50 %	VITAMIN A	IVIIIN	4000 IU/LB

INGREDIENTS

PROCESSED GRAIN BY-PRODUCTS, PLANT PROTEIN PRODUCTS, GRAIN PRODUCTS, MOLASSES PRODUCTS GROUND LIMESTONE, ROUGHAGE PRODUCTS (15%), SALT, DRIED ASPERGILLUS ORYZAE FERMENTATION PRODUCT, YEAST CULTURE, HYDROLYZED YEAST, VITAMIN A SUPPLEMENT, VITAMIN D3 SUPPLEMENT, VITAMIN E SUPPLEMENT, ZINC PROTEINATE, COPPER PROTEINATE, MANGANESE PROTEINATE, COBALT PROTEINATE, ZINC SULFATE, CALCIUM CARBONATE, FERROUS CARBONATE, MANGANESE SULFATE, COPPER SULFATE, SODIUM SELENITE, ETHYLENEDIAMINE DIHYDRIODIDE, COBALT SULFATE.

FEEDING DIRECTIONS

FEED CONTIUNALLY AT A RATE OF 10-40 MG/HEAD/DAY IN A LEAST 1 POUND AND NOT MORE THAN 10 POUNDS OF FEED. DAILY BAMBERMYCINS INTAKE IN EXCESS OF 20 MG/HEAD/DAY HAVE NOT BEEN SHOWN TO BE MORE EFFECTIVE THAN 20 MG/HEAD/DAY.

CAUTION: CONTAINS COPPER: COPPER HAS BEEN SHOWN TO BE TOXIC TO SHEEP.

3/8

ALLIED FEEDS INC. 208 HUTCHESON, CUERO, TEXAS 77954 1-800-242-3500 NET WT 50# (22.72 KG) 01



