

7130-BL

12% BEEF GROWER

MEDICATED

FOR BEEF STOCKER CALVES

For increased rate of weight gain in pasture cattle (slaughter, stocker feeder cattle, and dairy and beef replacement heifers)

ACTIVE DRUG INGREDIENTS

BAMBERMYCINS	4	GM/TC	NC
--------------	---	-------	----

GUARANTEED AMALTSIS

Commentation and the Comment of the				
CRUDE PROTEIN MIN 12.00 %	PHOSPHORUSMIN 0.50 %			
(This includes not more than 3.0% equi-	SALTMIN 1.50 %			
valent protein from non-protein nitroge	n.) SALTMAX 2.50 %			
CRUDE FAT MIN 3.40 %	POTASSIUMMIN 0.60 %			
CRUDE FIBER MAX 16.00 %	VITAMIN AMIN 5000 IU/LB			
CALCIUM MIN 0.95 %				
CALCIUM MAX 1.45 %				

MGREDIENTS

PROCESSED GRAIN BY-PRODUCTS, GRAIN PRODUCTS, ROUGHAGE PRODUCTS(18%), PLANT PROTEIN PRODUCTS, GROUND LIMESTONE, MOLASSES PRODUCTS, SALT, UREA, VITAMIN A SUPPLEMENT, D-ACTIVATED ANIMAL STEROL (SOURCE OF VITAMIN D3), VITAMIN E SUPPLEMENT, ZINC SULFATE, ZINC PROTEINATE, ASPERGILLUS ORYZAE FERMENTATION PRODUCT, YEAST CULTURE, HYDROLYZED YEAST, CALCIUM CARBONATE, FERROUS CARBONATE, MANGANESE SULFATE, MANGANESE PROTEINATE, COPPER SULFATE, COPPER PROTEINATE, SODIUM SELENITE, ETHYLENEDIAMINE DIHYDRIODIDE, COBALT SULFATE, COBALT PROTEINATE.

Feeding Directions

FEED CONTINUOUSLY AT A RATE OF 10 –40 MG/HEAD/DAY IN AT LEAST 1 POUND AND NOT MORE THAN 10 POUNDS OF FEED. DAILY BAMBERMYCINS INTAKE IN EXCESS OF 20 MG/HEAD/DAY HAVE NOT BEEN SHOWN TO MORE EFFECTIVE THAN 20 MG/HEAD/DAY.

3/8"

ALLIED FEEDS INC. 208 HUTCHESON , CUERO, TEXAS 77954 Totes or NET WT 50# (22.72 KG) 1-800-242-3500 01

