

## 9250-DE

# **14% DOE RATION**

## MEDICATED FOR RUMINANTS ONLY

## FOR GROWING AND FINISHING MEAT GOATS FOR SHOW

For the prevention of coccidiosis in young goats caused by Eimeria christenseni & Eimeria ninakohlykimovae

## ACTIVE DRUG INGREDIENTS

Decoquinate	13.6	g/te	on
-------------	------	------	----

#### GUARANTEED ANALYSIS

CRUDE PROTEINMIN	14.00 %	SALTMIN	1.00 %
This includes not more than		SALTMAX	1.25 %
1.0% equivalent crude protein		COPPERPPM	26.00
from non-protein nitrogen.		COPPERPPM	32.00
CRUDE FATMIN	2.50 %	MANGANESEPPM	90.00
CRUDE FIBERMAX	17.00 %	ZINCPPM	100.00
CALCIUMMIN	1.00 %	SELENIUMPPM	0.40
CALCIUMMAX	1.25 %	VITAMIN A .NOT LESS THAN IU/LB	3900
PHOSPHORUSMIN	0.5 %	VITAMIN E .NOT LESS THAN IU/LB	20

### INGREDIENTS

PROCESSED GRAIN BY-PRODUCTS, ROUGHAGE PRODUCTS (27%), GRAIN PRODUCTS, COTTON-SEED MEAL, MOLASSES PRODUCTS, GROUND LIMESTONE, SALT, AMMONIUM CHLORIDE, YEAST CULTURE, VITAMIN A SUPPLEMENT, VITAMIN D3 SUPPLEMENT, VITAMIN E SUPPLEMENT, MANGANESE SULFATE, MANGANESE PROTEINATE, FERROUS CARBONATE, ETHYLENEDIAMINE DIHYDRIODIDE, ZINC SULFATE, ZINC PROTEINATE, COPPER SULFATE, COPPER PROTEINATE, COBALT SULFATE, FERROUS SULFATE, COBALT PROTEINATE, SODIUM SELENITE.

WARNING: Do not feed to goats producing milk for human consumption.

### FEEDING DIRECTIONS

FEEDING 1# OF 14% DOE RATION SUPPLIES 6.8 mg OF DECOQUINATE. FEED AT LEAST 28 DAYS DURING EXPOSURE TO COCCIDIOSIS OR WHEN EXPERIENCE INDICATES THAT IT IS LIKELY TO BE A HAZARD.

PROVIDE FREE CHOICE MINERALS AND FRESH WATER AT ALL TIMES.

ALLIED FEEDS INC.

208 HUTCHESON : CUERO, TEXAS 77954

NET WEIGHT 50# (22.72 kg)